

Connecting you to the support you deserve

Indication

NUPLAZID® (pimavanserin) is a prescription medicine used to treat hallucinations and delusions associated with Parkinson's disease psychosis.

Important Safety Information for NUPLAZID (pimavanserin)

WARNING: INCREASED RISK OF DEATH IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

- Medicines like NUPLAZID can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia).
- NUPLAZID is not approved for the treatment of patients with dementia-related psychosis unrelated to the hallucinations and delusions associated with Parkinson's disease psychosis.

See additional Important Safety Information on back cover.
Please read the accompanying full Prescribing Information,
including **Boxed WARNING**, also available at NUPLAZID.com.

ONCE-DAILY
NUPLAZID[®]
(pimavanserin) 34mg capsules

A clear path forward

Beginning a new medication can feel overwhelming and confusing for you and your loved ones. NUPLAZIDconnect™ was specifically created to help you get the support you deserve for a clear path forward.

Your dedicated NUPLAZIDconnect Care Coordinator will connect you with the answers you need so you have a clear understanding of what to expect.

“[NUPLAZIDconnect] answered questions that I didn’t even know to ask. I cried—this was exactly what I needed.

CAREGIVER & DAUGHTER OF A LOVED ONE
LIVING WITH HALLUCINATIONS AND DELUSIONS
ASSOCIATED WITH PARKINSON’S DISEASE

Actor Portrayal

Your Care Coordinator can help you in a variety of ways, including:



Navigating insurance and managing costs

Your Care Coordinator can help explain your insurance coverage and the financial support options available to you.



Getting your prescription filled

Your Care Coordinator will work with your specialty pharmacy to fill your NUPLAZID (pimavanserin) prescription. They’ll make sure you understand when and how your medication will be delivered to you.



Answering your questions

Your Care Coordinator can also connect you to a Nurse Educator to answer questions you may have about your medication, such as:

- When and how to take NUPLAZID
- What side effects you may experience
- How long it may take for NUPLAZID to start working

Please don’t hesitate to reach out—we’re here for you and look forward to helping you find a clear path forward with NUPLAZIDconnect



Call 1-844-737-2223, Monday through Friday 8:30 AM–8:30 PM EST,
or visit [NUPLAZIDconnect.com](https://www.nuplazidconnect.com)

See Important Safety Information, including **Boxed WARNING**, on back cover. Please read the accompanying full Prescribing Information, also available at [NUPLAZID.com](https://www.nuplazid.com).

ONCE-DAILY
NUPLAZID
(pimavanserin) 34mg capsules

Important Safety Information and Indication for NUPLAZID® (pimavanserin) (cont.)

WARNING: INCREASED RISK OF DEATH IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

- Medicines like NUPLAZID can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia).
- NUPLAZID is not approved for the treatment of patients with dementia-related psychosis unrelated to the hallucinations and delusions associated with Parkinson's disease psychosis.
- **Do not take NUPLAZID if you** have had an allergic reaction to any of the ingredients in NUPLAZID. Allergic reactions have included rash, hives, swelling of the tongue, mouth, lips, or face, throat tightness, and shortness of breath.
- **NUPLAZID may cause serious side effects including:**
 - **QT Interval Prolongation:** NUPLAZID may increase the risk of changes to your heart rhythm. This risk may increase if NUPLAZID is taken with certain other medications known to prolong the QT interval. **Tell your healthcare provider about all the medicines you take or have recently taken.**
 - Do not take NUPLAZID if you have certain heart conditions that change your heart rhythm. It is important to talk to your healthcare provider about this possible side effect. Call your healthcare provider if you feel a change in your heartbeat.
- **Tell your healthcare provider about all the medicines you take.** Other medicines may affect how NUPLAZID works. Some medicines should not be taken with NUPLAZID. Your healthcare provider can tell you if it is safe to take NUPLAZID with your other medicines. Do not start or stop any medicines while taking NUPLAZID without talking to your healthcare provider first.
- The **most common side effects** of NUPLAZID include swelling in the legs or arms, nausea, confusion, hallucination, constipation, and changes to normal walking. These are not all the possible side effects of NUPLAZID. For more information, ask your healthcare provider about this medicine.

Indication

NUPLAZID (pimavanserin) is a prescription medicine used to treat hallucinations and delusions associated with Parkinson's disease psychosis.

Dosage and Administration

The recommended dose of NUPLAZID is one 34 mg capsule once per day, taken by mouth.

NUPLAZID is available as 34 mg capsules and 10 mg tablets.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You can also call ACADIA Pharmaceuticals Inc. at 1-844-4ACADIA (1-844-422-2342).

Please read the accompanying Prescribing Information, including **Boxed WARNING**, also available at NUPLAZID.com.

©2019 ACADIA Pharmaceuticals Inc. ACADIA, NUPLAZID, and NUPLAZIDconnect are trademarks of ACADIA Pharmaceuticals Inc. All rights reserved. NC-0131-v2 07/19.